



Statistics &amp; Probability Letters 47 (2000) 209–211

---

**STATISTICS &  
PROBABILITY  
LETTERS**


---

[www.elsevier.nl/locate/stapro](http://www.elsevier.nl/locate/stapro)

## Addendum

# Addendum to “A note on complete convergence for arrays”, Statist. Probab. Lett. 38 (1) (1998) 27–31<sup>☆</sup>

Tien-Chung Hu<sup>a,\*</sup>, Andrei Volodin<sup>b</sup><sup>a</sup>*Department of Mathematics, National Tsing Hua University, Hsinchu 300, Taiwan*<sup>b</sup>*Research Institute of Math. and Mech., Kazan University, Kazan 420008, Russia*

Received August 1999

**Abstract**

Under some conditions on an array of rowwise-independent random variables, Hu, Szynal and Volodin obtained a complete convergence result for law of large numbers. In this addendum we mention that the convergent rate of sequence  $\{c_n, n \geq 1\}$  must be bounded away from zero. © 2000 Elsevier Science B.V. All rights reserved

MSC: 60F15; 60G50

**Keywords:** Arrays; Rowwise independence; Sums of independent random variables; Complete convergence; Weak law of large numbers

In our paper (Hu et al., 1998), the following complete convergence theorem for arrays of rowwise-independent random variables was formulated.

**Theorem.** Let  $\{X_{ni}, 1 \leq i \leq k_n, n \geq 1\}$  be an array of rowwise-independent random variables and  $\{c_n, n \geq 1\}$  be a sequence of positive constants such that

$$\sum_{n=1}^{\infty} c_n = \infty. \quad (1)$$

Suppose that for every  $\varepsilon > 0$  and some  $\delta > 0$ :

- (i)  $\sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} P(|X_{ni}| > \varepsilon) < \infty,$

---

<sup>☆</sup> PII of the original article: S0167-7152(98)00150-8.

\* Corresponding author.

E-mail address: [tchu@math.nthu.edu.tw](mailto:tchu@math.nthu.edu.tw) (T.-C. Hu)